

Developing Attract-and-kill Strategies To Manage Spotted Wing Drosophila, *Drosophila Suzukii* Matsumara, In Raspberry.




Male

Spotted Wing Drosophila



Female

Peter Jentsch
Extension Associate



Cornell University

Hudson Valley Research Laboratory


Questions and Objectives

- What components offer effective olfactory, visual and sensory perception for attraction?
- Can we construct an long lasting, economically viable Attract and Kill station to reduce SWD populations leading to reduced fruit injury?
- Can Attract and Kill (AtK) Technologies work well alone or do they require the combination of other management strategies to reduce pesticide loading in small fruit production systems?



SWD Spread from 2008 – 2013 in the US

Hawaii 1980




Cornell University

Hudson Valley Research Laboratory


SWD in New England - 2016

First NY detection 2011 in NY
Mid-September Organic Raspberry
Rensselaer & Suffolk Counties ●



SWD in New York - 2016

First State Capture ●
June 21, 2016
Wayne County, NY



Cornell University

Hudson Valley Research Laboratory

Success of SWD in Small Fruit




Occupies a relatively non-competitive niche

- Able to penetrate and oviposit into un-ripened fruit using a highly sclerotized & serrated ovipositor .



Reproductive Success of SWD in Small Fruit



- Optimal development is at 65-70°F, ~**12 day** generation time.
- Adult flies live for 3-6 weeks, and females can lay over 300 eggs.
- Limited by high heat in summer and by winter cold. But, SWD populations are found in cold regions of Japan.
- 3-10 generations in NY



Fruit Affected by SWD

Highest risk

Strawberries

Raspberries

Cherries (Late var,)

Nectarines

Blueberries

Blackberries

Moderate risk

Peaches

Grapes

Pears

Apples

Tomato

Alternate hosts

Wild plants with berries,
such as...



Tartarian Honeysuckle

Snowberry

Elderberry

Pokeweed

Dogwood



Cornell University

Hudson Valley Research Laboratory

SWD Alternate Host: Population Development in the HV

Monitoring *L. tartarica*



- Honeysuckle is a primary host for SWD; *L. tartarica* fruit favored over raspberry in June-August.



Sampling and Monitoring Protocols

Monitoring


Weekly trap captures, 4 traps per site including Tartarian Honeysuckle *L. tartarica*

Extension Outreach: EDDMaps

Sampling

Weekly 25 fruit from each of 4 plant clusters (10')







Weigh and assess fruit for SWD eggs (expressed as eggs/gram)



Cornell University

Hudson Valley Research Laboratory

Monitoring SWD Using ACV on 6 Farms in the Hudson Valley Eastern, NY - 2012



Fruit Monitoring & Injury

- SWD oviposition may precedes adult trap captures in production systems.
- Newer traps have increased sensitivity to adult presence
- Conventional and organic production systems contain raspberry fruit with SWD eggs & larva.



Managing Insecticide Resistance: Raspberry




SWD Control in Mixed Small Fruit; Orange Co. 2012

Date	Material	Rate	Commodity
27 June	Malathion 57	2 pts./A	Raspberry
1 July	Assail 30SG	5 oz./A	Raspberry
5 July	Malathion 57	2 pts./A	Raspberry
12 July	Delegate 25WDG	3 oz./A	Raspberry
14 July	Brigade	8 oz./A	Raspberry
19 July	Assail 30SG	5 oz./A	Raspberry
22 July	Danitol	16 oz./A	Raspberry
27 July	Mustang Max	4 oz./A	Raspberry
30 July	Assail 30SG	5 oz./A	Raspberry
6.31" Rainfall; 6 day application interval			
5 August	Delegate 25WDG	3 oz./A	Raspberry
19 August	Brigade	8 oz./A	Raspberry



Managing Insecticide Resistance: Raspberry



AtK Based Literature Eastern US

Tracy Leskey (USDA-ARS)

Developing a Behaviorally Based Attract and Kill System for SWD

- **Color important;** black and red routinely outperformed other colors.
- **A spherical shape: size greater than 2.5 cm acceptable.**
- **Baits** enhance SWD capture
- **SWD infestation in raspberries reduced by 50% when sphere with sugar and bait in caged studies. Sprayed fruit + AtK in combination most effective in managing SWD compared to either alone under high pressure.**



Cesar Rodriguez-Saona, Rutgers State U. Of N.J., Richard Cowles Univ. Conn.

Bait comparisons of SWD in blueberry

- **Suzukii and Trece baits very effective at capturing SWD with Trece and apple cider vinegar capturing higher numbers of non-SWD flies.**

Cowles, R. S., C. Rodriguez-Saona, R. Holdcraft, G. M. Loeb, J. E. Elsensohn, and S. P. Hesler. 2015. Sucrose improves insecticide activity against *Drosophila suzukii* (Diptera: Drosophilidae). J. Econ. Entomol. 1 – 14. DOI: 10.1093/jee/tou100.



AtK Based Literature Western US



Alan L Knight, Esteban Basoalto, Wee Yee. **Adding Yeasts with Sugar** to Increase the Number of Effective Insecticide Classes to Manage *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae) in Cherry Pest Management Science · October 2015

Alan L Knight, Esteban Basoalto, Wee Yee. Developing a new bait for spotted-wing drosophila in organic cherry production Acta horticulturae 1001(1001):147-152 · July 2013

Increased attractiveness of bait using bread yeast, *Saccharomyces cerevisiae*

- **Exceeds the attractiveness of commercial products GF-120® and Nu-Lure®,**
- **Addition of the sugar-yeast bait to Entrust increased fly mortality 4-fold in early-season bioassays with green and yellow cherries, reducing eggs laid and larval infestations by 50%**




SWD Adult Preference Binary Choice Tests

Mean # AtK Component Attractiveness



■ # Male SWD ■ # Female SWD ■ # Total

25% Red Raspberry Concentrate
75% Apple Cider Vinegar
Brewers yeast: *Saccharomyces cerevisiae*



Cornell University

Hudson Valley Research Laboratory

Methods: Development of Attract and Kill for Management of SWD in Small Fruit



AtK Construction



- 3" substrate woven polypropylene netting as a base
- Super Absorbent Polymer (SAP)
- Gelatin
- Red raspberry concentrate
- Apple cider vinegar
- Brewers yeast
- 1% A.I.
- AtK solution applied at 2 mL/disk



Methods: Development of Attract and Kill for Management of SWD in Small Fruit




Insecticide Product	Active Ingredient (IRAC Group)
Malathion 5EC	malathion (IRAC 1B)
Imidan 70W	phosmet (IRAC 1B)
Assail 30SG	acetamiprid (IRAC 4A)
Scorpion 35 SL	dinotefuran (IRAC 4A)
Brigade EC	bifenthrin (IRAC 3A)
Mutang Max	zeta-cypermethrin (IRAC 3A)
Pyganic EC 1.4	pyrethrin (IRAC 3A)
Triple Crown	bifenthrin, imidacloprid, zeta-cypermethrin (IRAC 3A, 4A)
Delegate WG	spinetoram (IRAC 5)
Entrust SC	spinosad (IRAC 5)
Exirel	cyazypyr (IRAC 28)
BotaniGard; Mycotrol	<i>Beauveria bassiana</i> strain GHA
BalEnce	<i>Beauveria bassiana</i> Diptera-specific strain (HF23)
Boric Acid	99% Boric Acid
Hot Shot Maxattrax Roach Powder	99% Boric Acid formulated




Attract and Kill Station Efficacy

Lab Caged Studies (25 SWD 48h 75F 75%rH 14/10 LD)



Attract and Kill Station Recharge Efficacy

**SWD Eggs Per Gram of Raspberry & Adult Mortality @ 72h
24h (Wet) vs 7d (Dry) treated disks**



1% A.I. Entrust (spinosad-Dow)




Cornell University

Hudson Valley Research Laboratory

Attract and Kill Station Recharge Efficacy



SWD Adult Mortality



1% A.I. Entrust (spinosad-Dow)




Cornell University

Hudson Valley Research Laboratory

Attract and Kill Station Recharge Efficacy



Eggs Per Gram in Raspberry Fruit




1% A.I. Entrust (spinosad-Dow)



Cornell University

Hudson Valley Research Laboratory

Insecticidal Options for AtK Stations




Observations

- Initial weight loss of $\geq 50\%$ in 30 hours and overall seasonal weight loss of 70%.
- Extended rain events increase fluctuations in AtK disk weight.



Attract and Kill Station Recharge Efficacy



Observations

- Extended high relative humidity also increase weight.
- Inversely, low rH reduces weight.
- Morning dew is also absorbed by the disk.



Attraction of *Drosophila* to AtK from Morning Dew




June 14th – September 19th 8:30 AM,



Cornell University

Hudson Valley Research Laboratory

Experimental Field Design*




3 Raspberry Plantings on 3 Farm sites in two NY counties 1 Conventional & 2 Organic Production Systems

AtK placement timed for each row (A,B,C)

- A. 1st SWD in NY (14th June)
- B. 1st SWD on site (19th June)
- C. 1st SWD oviposition of fruit (25th June)

* Row spacing- 11'; plant spacing 3'; 2 of 3 sites used wire trellis used to hang AtK stations

Experimental Field Design



3 Raspberry Plantings on 3 Farm sites in two NY counties 1 Conventional & 2 Organic Production Systems

AtK placement timed for each row (A,B,C)

- A. 1st SWD in NY (14th June)
- B. 1st SWD on site (19th June)
- C. 1st SWD oviposition of fruit (25th June)

Split Block


(Reps I-III)

Red and Yellow Disk sprayed weekly

(Reps IV-VI)

Red and Yellow Disk sprayed 2x/week

Experimental Field Design



3 Raspberry Plantings on 3 Farm sites in two NY counties 1 Conventional & 2 Organic Production Systems

AtK placement timed for each row (A,B,C)

- A. 1st SWD in NY (14th June)
- B. 1st SWD on site (19th June)
- C. 1st SWD oviposition of fruit (25th June)

Split Block




(Reps I-III)

Red and Yellow Disk sprayed weekly


(Reps IV-VI)

Red and Yellow Disk sprayed 2x/week

Treatments

Red		1% Borax treated disks spaced at 1.5' (120) Disks/ side = 240 disks/ row
Yellow		1% Borax treated disks spaced at 3' (60) Disks/ side = 120 disks/ row
Green		Untreated disks spaced at 3' (60) Disks/ side = 120 disks/ row

AtK Treatment Spacing




SWD Damage Means in Raspberry Fruit

Atk Management of SWD in Raspberry Trapanni Orchard, Marlboro, NY - 2016

F-Value
0.99051


P-Value
0.4415



SWD Damage Means in Raspberry Fruit

AtK Management of SWD in Raspberry WestWind Orchard, Accord , NY - 2016


F-Value 0.53805
P-Value 0.7993



SWD Damage Means in Raspberry Fruit

AtK Management of SWD in Raspberry
PFP Organic CSA, Poughkeepsie , NY - 2016


F-Value 7.02602
P-Value 0.0001



Combined Farm & AtK Application Timing

AtK Management of SWD in Raspberry Fruit Damage Means, Ulster & Dutchess Co; NY - 2016

F-Value 1.64091
P-Value 0.1252



*59.9% Reduction of Raspberry Fruit Injury over the UTC



Cornell University

Hudson Valley Research Laboratory

Conclusion



- **Attract and kill strategies have been shown to provide reduced levels of infestation from spotted wing drosophila in conventional and organic raspberry production systems.**
- **Further study of placement density and reapplication intervals of AtK disks for optimal control is needed prior to recommendations for use.**
- **Use of AtK + 1% Boric Acid in combination with cultural control, frequent harvest intervals, berry sanitation and harvest low temperature storage strategies may decrease the impact of SWD while reducing the resistance potential in SWD populations from frequent insecticide use.**



Partnership Thanks

- New York Farm Viability Grant - OAR 15 013
- Greg Loab, NYSAES, Geneva, NY
- Juliet Carrol, NYS IPM, Geneva, NY
- Tim Lampasona, Jonathon Binder, Mike Fraatz
Hudson Valley Research Laboratory

Fabio Chizola, WestWind Farm, Accord, NY

Poughkeepsie Farm Project, Poughkeepsie, NY

Trapani Farm & Orchard, Marlboro, NY



WESTWIND
ORCHARD



Cornell University

Hudson Valley Research Laboratory