

Reprint Series
16 March 1990, Volume 247, pp. 1325–1328

SCIENCE

**Fossil Soils and Grasses of a Middle Miocene
East African Grassland**

GREGORY J. RETALLACK, D. P. DUGAS, AND E. A. BESTLAND

Fossil Soils and Grasses of a Middle Miocene East African Grassland

GREGORY J. RETALLACK, D. P. DUGAS, E. A. BESTLAND

Fossil soils and grasses from the well-known Miocene mammal locality of Fort Ternan, southwestern Kenya, are evidence of a mosaic of grassy woodland and wooded grassland some 14 million years ago. This most ancient wooded grassland yet known on the African continent supported more abundant and diverse antelopes than known earlier in Africa. Ape fossils at Fort Ternan, including *Kenyapithecus wickeri*, were associated with woodland parts of the vegetation mosaic revealed by paleosols. Grassland habitats were available in East Africa long before the evolutionary divergence of apes and humans some 5 to 10 million years ago.

SAVANNA ECOSYSTEMS HAVE LONG been linked with early human evolution, and the antiquity of savannas in Africa has been a source of debate (1). Part of the problem is the loose use of the term "savanna" to include any kind of tropical grassland. We prefer the term "wooded grassland" for well-drained grassy vegetation with 10 to 40% cover by trees (2). Unlike wooded and open grassland, grassy woodland and marsh are at least as old as 23 million years and remained widespread 17 million years ago in East Africa, judging from evidence of paleosols (3), of fossil sedges and grasses (4), of fossil dicots (5), and of fossil birds allied to modern forms now found widely in grassy vegetation (6). Of more interest from the point of view of the evolution of African mammals is the origin of wooded grassland and open grassland among more ancient kinds of woodland and forest vegetation.

Paleosols can be useful in deciphering ancient vegetation mosaics, because, unlike fossils, they are by definition in the place they formed. They are also abundant and have been widely recognized in southwest

Kenya (3, 7-9). The thin, brown, nodular, calcareous paleosols found throughout the 9.6-km outcrop of the middle Miocene Fort Ternan Beds are distinct from geologically older paleosols in Kenya (3, 10). The Fort Ternan beds are a sequence of carbonatite-nephelinite tuffs, lahars, colluvium, and alluvium. They have been dated by the K/Ar method at 14.4 ± 0.2 million years old by the use of biotite (7, 11). This account details only paleosols in the 8-m-thick sequence exposed in the large (12) quarry for fossils at Fort Ternan National Monument. Three distinct kinds of paleosols were recognized and named informally from the local Dholuo language: Chogo ("bone"), Onuria ("yellow"), and Dhero ("thin") paleosol series. The paleosols were characterized in the field, petrographically, and chemically (13). From these data (Fig. 1), some aspects of the original soils and their environment can be reconstructed.

One approach to interpreting paleosols is to identify them within a soil classification and compare them to modern soils, with allowance for possible alteration after burial. The distinctive surface horizons of both Onuria and Chogo paleosols are critical to their identification: they have the granular structure, thickness, dark color, and large proportion of bases that define a mollic epipedon and Mollisols (14). Organic mat-

G. J. Retallack and E. A. Bestland, Department of Geological Sciences, University of Oregon, Eugene, OR 97403.

D. P. Dugas, Department of Geography, University of Oregon, Eugene, OR 97403.

ter is not sufficiently abundant to meet mollic criteria, but it is known that organic matter is lost by about an order of magnitude from well-drained paleosols soon after burial (15). A mollic interpretation is not compromised either by an early burial cement of un-recrystallized sparry calcite in sandy layers or compaction of clayey parts of the profiles to an estimated 74% of their former thickness (16). The Chogo clays can be identified as Haplustolls and the Onuria clay as a Calcustoll (14). Both Chogo and Onuria paleosols are similar in profile form and in chemical and petrographic depth functions to modern soils formed on carbonatite-nephelinite ash under mid-length grassland and wooded grassland in mid-slope positions in the central Serengeti Plains of Tanzania (17, 18). They are neither so clayey, deeply weathered, or red as African volcanic soils under lowland or montane woodland or forest (18, 19) or geologically older paleosols in Kenya (3, 9, 10), nor are they so thin, zeolitic, or calcareous as East

African desert soils on carbonatite-nephelinite ash (20).

A second approach for interpreting these paleosols is to assess soil features under the familiar headings of climate, organisms, topographic relief, parent material, and time for formation. The length of time represented by calcareous paleosols can be assessed from the morphology of their subsurface calcareous (Bk, Fig. 1) horizons (21, 22). Both Chogo and Onuria paleosols have distinct nodules with replacive and displacive microfabrics. In contrast, Dhero paleosols show little discernible pedogenic reorganization of carbonate. Like similar surface soils of known age in the Serengeti Plains of Tanzania (21), Chogo and Onuria paleosols may represent 2000 to 9000 years of soil formation; and Dhero paleosols, a few hundred years or less.

Topographic setting and parent materials for the paleosols are known from geological context to be on the lowest slopes of a carbonatite-nephelinite volcano (7, 8). On-

uria and Dhero series paleosols formed on lahars; and Chogo paleosols on colluvium. These indications of good drainage and base-rich parent materials are compatible with the lack of petrographic or chemical evidence for salinization, gleization, or leaching in all the paleosols (Fig. 1). Thus, these paleosols were not restricted to salt licks, seasonally waterlogged depressions (dambos), or marshes.

Indications of grassy vegetation are provided by abundant fine root traces in all the paleosols. Fossil stumps are rooted in Dhero paleosols. These paleosols may have supported early successional vegetation, unlike Chogo and Onuria paleosols with little relict bedding. Two small stump casts 7.5 and 16 cm in diameter and 2 m apart were seen in the Chogo clay eroded-phase paleosol 5 m west along strike from the type Chogo clay. This paleosol with fossil stumps and large root traces also has an incipient subsurface clayey horizon and diffuse boundaries between horizons, as in the Chogo clay ferru-




Fig. 1. Detailed sections of the four different kinds of paleosols recognized within the 8-m sequence exposed in the main fossil excavation at Fort Ternan National Monument, Kenya (13). Heights in centimeters are calibrated to a

longer section in which the top of the underlying Baraget Phonolite is at 780 cm.

ginized nodule variant (Fig. 1) and modern soils of grassy woodland (18). Paleosols lacking stumps, such as the Onuria and type Chogo clay paleosols, have an abrupt and planar lower boundary to the dark surface (A, Fig. 1) horizon, as in modern soils of tall to intermediate grassland and wooded grassland (17). This evidence for a local vegetation mosaic in a 7-m lateral exposure of Chogo paleosols is not apparent from scattered fossil fruits and seeds of woodland trees and vines or leaves and pollen of grasses, recovered from Chogo paleosols (23). Abundant bones were fossilized in place as an attritional assemblage in Chogo paleosols (7). The most common large mammals were antelope (*Oioceros* and *Kipsigicerus*), giraffoid (*Palaeotragus*), rhinoceros (*Paradiceros*), and proboscidean (*Choerolophodon*). Rhinos and proboscideans had a prior evolutionary history in East Africa (24), but the antelopes were newly immigrant to this area. In tooth and limb morphology, the antelopes were better suited to open country than were early Miocene African mammals but in these adaptations were most like modern woodland antelope (25). Fossil snails from Chogo paleosols reflect the wooded part of the mosaic (26).

Mean annual rainfall at Fort Ternan was probably in the semiarid range, because the paleosols would have had a calcareous horizon at 33 cm below the surface before burial compaction (16), as in moderately developed soils of the central Serengeti Plains receiving 320 to 590 mm of rainfall (17). Similar estimates of former rainfall are gained by comparison with moderately developed soils in the United States and India (27).

Additional indications of tropical grassland at Fort Ternan are fossil grasses found in several of the paleosols, but best preserved in calcareous nephelinitic sandstone covering the Onuria paleosol (28). The grasses are strongly oriented as if pushed over in place of growth by flow from the north (29). In the 7-m length of the grass bed exposed in the quarry, grasses are confined to its basal 17 cm with little clumping. Many of the grasses were stout and long: some were seen up to 8 mm in diameter and 24 cm long. From these observations, the grass bed has preserved a sod-forming, mid-length to tall grassy sward.

With the scanning electron microscope (SEM), a few small areas of some fossil grasses showed well preserved stomates and phytoliths (Fig. 2). The best results were obtained with specimens mounted on stubs and coated immediately after extraction from the matrix. Cuticular details of the grasses were compared with those from SEM studies of living East African grasses




Fig. 2. Miocene fossil grasses from Fort Ternan (28): including a branching panicoid grass (A) and one of its immature stomates with domed subsidiary cells from a basal leaf blade (C); a second species of panicoid grass with abundant dumbbell-shaped and common three-lobed silica bodies (B), and a chloridoid grass with circular silica bodies (D). Specimens and SEM stubs are from the Kenyan National Museum, Nairobi, numbers FT-13126 (A and C), FT-13120 (B), and FT-13127 (D).

(30) and were classified to subfamily level by reference to recent compilations of grass epidermal features (31). Five distinctly different morphologies and arrangements of stomates, phytoliths, and hair bases were found among the fossil grasses. Three of them are identified as different grass species of the subfamily Panicoidae and two as separate species of the subfamily Chloridoideae (28). Modern chloridoid species most like the fossils in cuticular morphology are found mostly in open areas prone to grazing and other disturbance. Modern panicoids most like the fossils live in grasslands and grassy woodlands. There were no bambusoid or arundinoid grasses, common in seasonally wet grasslands (dambos), woodlands, or forests (2, 32), nor were there pooid grasses that now dominate Afrotropical and temperate vegetation (2, 31).

These paleoenvironmental inferences are confirmed by the functional morphology of the fossil cuticles. These were not arid or perarid grasslands, because the fossil stomates are not deeply sunken or hidden within stomatal grooves (33). Indications of high grazing pressure on the grasses are provided by the abundance of phytoliths on the fossil cuticles, including highly abrasive dumbbell and three-lobed forms (34). In both subfamilial taxonomic affinities and in their functional morphology, the fossil grasses compare well with those of intermediate to tall grasslands and wooded grasslands of the modern Serengeti Plains of Tanzania (2, 17).

Fossil apes from Fort Ternan, including *Kenyapithecus wickeri* (8, 26), have been recorded in attritional assemblages in place (7) only in the Chogo clay eroded-phase paleosol, here interpreted to have formed under grassy woodland at an ecotone to wooded grassland on the adjacent type Chogo clay. Such a habitat for *K. wickeri* is compatible with what little is known about its limb structure (35). The evidence presented here brings into clearer focus the array of potential environments for human evolution from apes, which may have occurred in East Africa 5 to 10 million years ago (1). Forests and woodlands certainly were present during middle and late Miocene time (3, 32) and have persisted to this day in well-watered areas. However, grasses and soils structurally and taxonomically similar to those of mosaics of grassy woodland and open wooded grassland also have been present for at least the past 14 million years in East Africa.

REFERENCES AND NOTES

1. A. Hill, *J. Hum. Evol.* **16**, 583 (1987).
2. F. White, *The Vegetation of Africa* (Unesco, Paris, 1983).
3. G. J. Retallack, *Soils of the Past* (Unwin-Hyman, London, 1990).
4. A. C. Hamilton, *Uganda J.* **32**, 157 (1968).
5. D. I. Axelrod and P. H. Raven, in *Biogeography and Ecology of Southern Africa*, M. J. A. Werger and A. C. Van Bruggen, Eds. (Junk, The Hague, 1978), pp. 77-130.
6. S. L. Olson and D. T. Rasmussen, *Science* **233**, 1202 (1986).
7. P. Shipman, A. Walker, J. A. Van Couvering, P. J.

- Hooker, J. A. Miller, *J. Hum. Evol.* **10**, 49 (1981).
8. M. Pickford, *Muench. Geowiss. Abh.* **8** (1986), 151 pp.
 9. G. D. Thackray, thesis, University of Oregon, Eugene (1989).
 10. Since 1984, our studies have characterized 95 paleosols of 20 kinds from Koru, Songhor, Rusinga Island, and Fort Ternan, ranging in geological age from 12 to 20 million years.
 11. Date corrected to new constants by the procedure of G. B. Dalrymple [*Geology (Boulder)* **7**, 558 (1979)].
 12. Floor area 132 m², wall length 46 m, and wall area about 165 m².
 13. Colors compared with a Munsell chart; grain size distribution and petrographic composition determined from 500 points in thin section with a Swift automatic point counter; G. S. Smith counted Onuria and Dhero paleosols of Fig. 1, and G.J.R. confirmed Onuria and counted Chogo paleosols; organic carbon by D. A. Horneck, Soil Testing Laboratory of Oregon State University, with the Walkley-Black technique; chemical analyses from Atomic Absorption by C. McBirney, University of Oregon.
 14. Terminology of Soil Survey Staff, *Handb. U.S. Dep. Agric.* **436** (1975).
 15. F. J. Stevenson, *Soil Sci.* **107**, 470 (1969).
 16. Burial by 600 m was estimated from a reconstruction of Tindereet volcano by analogy with the petrologically similar nearby volcano Kisingiri, studied by M. J. Le Bas [*Carbonatite-Nephelinite Volcanism* (Wiley, London, 1977)] and compaction from standard curve of B. Baldwin and C. O. Butler [*Bull. Am. Assoc. Petrol. Geol.* **69**, 622 (1985)].
 17. H. A. de Wit, *Soils and Grassland Types of the Serengeti Plain (Tanzania)*, (Mededelingen Landbouwhogeschool Wageningen, the Netherlands, 1978); G. D. Anderson and L. W. Talbot, *J. Ecol.* **53**, 33 (1965).
 18. T. J. Jager, *Soils of the Serengeti Woodlands, Tanzania* (Center for Agricultural Publishing and Documentation, Wageningen, the Netherlands, 1982).
 19. C. Mizota, I. Kawasaki, T. Wakatsuki, *Geoderma* **43**, 131 (1988).
 20. G. D. Anderson, *Sols Africains* **8**, 339 (1963); R. L. Hay, *J. Volcanol. Geotherm. Res.* **37**, 77 (1989).
 21. R. L. Hay and R. J. Reeder, *Sedimentology* **25**, 649 (1978).
 22. M. N. Machette, *Spec. Pap. Geol. Soc. Am.* **302** (1985), p. 1.
 23. P. L. Shipman, thesis, New York University (1977); R. Bonnefille, in *The Evolution of the East Asian Environment*, R. O. Whyte, Ed. (University of Hong Kong, Hong Kong, 1984), vol. 2, pp. 579-612.
 24. D. A. Hooijer, *Zool. Meded. (Leiden)* **43**, 77 (1968); P. Tassy, *Bull. Soc. Geol. Fr.* **21**, 265 (1979).
 25. H. Thomas, *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **40**, 251 (1984); J. Kappelman, *Am. J. Phys. Anthropol.* **75**, 229 (1988).
 26. M. Pickford, *J. Hum. Evol.* **14**, 113 (1985); *ibid.* **16**, 305 (1987).
 27. H. Jenny, *Factors in Soil Formation* (McGraw-Hill, New York, 1941); R. J. Arkley, *Soil Sci.* **96**, 239 (1963); J. L. Sehgal, C. Sys, D. R. Bumbala, *Pedologie* **18**, 351 (1968).
 28. D. P. Dugas, thesis, University of Oregon, Eugene (1989).
 29. A set of 50 fossil grasses excavated in the laboratory from a single large oriented sandstone block had a vector mean orientation of 351° azimuth and a vector magnitude of 82% (a highly consistent orientation). Another set of 35 grasses measured in the field had a vector mean of 349° and a vector magnitude of 90%. In both sets the grasses dipped consistently to the north.
 30. P. G. Palmer and A. E. Tucker, *Smithson. Contrib. Bot.* **49**, 1 (1981); *ibid.* **59**, 1 (1983); P. G. Palmer and S. Gerbeth-Jones, *ibid.* **62**, 1 (1986); *ibid.* **67**, 1 (1987); P. G. Palmer, S. Gerbeth-Jones, S. Hutchinson, *ibid.* **55**, 1 (1985).
 31. R. P. Ellis, in *Grass Systematics and Evolution*, T. R. Soderstrom, K. W. Hilu, C. S. Campbell, M. E. Barkworth, Eds. (Smithsonian Institution Press, Washington, DC, 1987), pp. 3-10.
 32. B. F. Jacobs and C. A. S. Kabuye, *J. Hum. Evol.* **16**, 147 (1987).
 33. C. R. Metcalfe, *Anatomy of the Monocotyledons* (Clarendon, Oxford, 1960).
 34. C. M. Herrera, *Oikos* **45**, 446 (1985).
 35. B. Senut, *Ann. Paleontol.* **74**, 128 (1988).
 36. P. Andrews, M. Pickford, P. Shipman, and A. Walker helped with discussion; fieldwork was authorized by the Office of the President of the Republic of Kenya, under the auspices of the National Museums of Kenya and their director, R. E. Leakey; funded by the Wenner Gren Foundation and NSF grant EAR 8503232 to G.J.R., and by a Condon Award of the University of Oregon and Geological Society of America grants 3753-87 and 3969-88 to D.P.D.

26 June 1989; accepted 30 January 1990

