




# Biomass Coverage in Pleuronectids Taxonomy Group




# Biomass Coverage Over Time in Pleuronectids Taxonomy Group

## Stocks with highest average biomass



# Biomass Coverage Over Time in Pleuronectids Taxonomy Group

## Stocks with lowest average biomass







Taxonomy Group

- Canada East Coast
- European Union (non Mediterranean)
- US East Coast
- Canada West Coast
- Northwest Pacific
- US West Coast




Summary method:  Median

 State-space model prediction




# Yellowfin sole Bering Sea and Aleutian Islands – Pleuronectids Taxonomy Group

Stock represents 16.1% of average absolute surplus production




# Arrowtooth flounder Gulf of Alaska – Pleuronectids Taxonomy Group

Stock represents 5.02% of average absolute surplus production




# Northern rock sole Eastern Bering Sea and Aleutian Islands – Pleuronectids Taxonomy Group

Stock represents 8.16% of average absolute surplus production



# Dover sole Pacific Coast – Pleuronectids Taxonomy Group


Stock represents 0.834% of average absolute surplus production








# Arrowtooth flounder Gulf of Alaska in Pleuronectids Taxonomy Group



First and last years with recorded surplus production are labeled

# Northern rock sole Eastern Bering Sea and Aleutian Islands in Pleuronectids Taxonomy Group



First and last years with recorded surplus production are labeled

# Dover sole Pacific Coast in Pleuronectids Taxonomy Group

